



**DEFENSE TECHNICAL INFORMATION CENTER**

*Information for the Defense Community*

DTIC® has determined on 11 / 18 / 13 that this Technical Document has the Distribution Statement checked below. The current distribution for this document can be found in the DTIC® Technical Report Database.

☒ **DISTRIBUTION STATEMENT A.** Approved for public release; distribution is unlimited.

☐ **© COPYRIGHTED.** U.S. Government or Federal Rights License. All other rights and uses except those permitted by copyright law are reserved by the copyright owner.

☐ **DISTRIBUTION STATEMENT B.** Distribution authorized to U.S. Government agencies only (fill in reason) (date of determination). Other requests for this document shall be referred to (insert controlling DoD office).

☐ **DISTRIBUTION STATEMENT C.** Distribution authorized to U.S. Government Agencies and their contractors (fill in reason) (date determination). Other requests for this document shall be referred to (insert controlling DoD office).

☐ **DISTRIBUTION STATEMENT D.** Distribution authorized to the Department of Defense and U.S. DoD contractors only (fill in reason) (date of determination). Other requests shall be referred to (insert controlling DoD office).

☐ **DISTRIBUTION STATEMENT E.** Distribution authorized to DoD Components only (fill in reason) (date of determination). Other requests shall be referred to (insert controlling DoD office).

☐ **DISTRIBUTION STATEMENT F.** Further dissemination only as directed by (insert controlling DoD office) (date of determination) or higher DoD authority.

*Distribution Statement F is also used when a document does not contain a distribution statement and no distribution statement can be determined.*

☐ **DISTRIBUTION STATEMENT X.** Distribution authorized to U.S. Government Agencies and private individuals or enterprises eligible to obtain export-controlled technical data in accordance with DoDD 5230.25; (date of determination). DoD Controlling Office is (insert controlling DoD office).

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
<b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</b>					
1. REPORT DATE (DD-MM-YYYY) 09/16/2008		2. REPORT TYPE Final		3. DATES COVERED (From - To) 06/15/2007 - 06/14/2008	
4. TITLE AND SUBTITLE DURIP: A Computational Cluster for Multiscale Simulations of Ionic Liquids				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER FA955007-1-0512	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Gregory A. Voth, PI				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Utah Department of Chemistry 315 S. 1400 E., Rm. 2020 Salt Lake City, UT 84112-0850				8. PERFORMING ORGANIZATION REPORT NUMBER  Final	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) University of Utah Office of Sponsored Projects 1471 E. Federal Way Salt Lake City, UT 84102-1821				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT N AFRL-OSR-VA-TR-2013-0960	
12. DISTRIBUTION/AVAILABILITY STATEMENT  <div style="text-align: center; font-size: 2em; margin-top: 20px;">20130918354</div>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The focus of this project was to acquire and use computer cluster nodes purchased with AFOSR DURIP funding for the multiscale simulation of ionic liquids and the construction of a novel molecular design methodology. During the funding period atomistic and coarse-grained (CG) molecular dynamics (MD) simulations of ionic liquids were performed to study the structure and dynamical properties of the ionic liquids. An effective force coarse graining (EF-CG) method was developed to build transferable CG models, and a systematic inverse coarse-graining molecular design approach was developed based on the EF-CG model and an empirical CG-structure-property relationship.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES  6	19a. NAME OF RESPONSIBLE PERSON Gregory A. Voth, PI
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code) 801-581-7272

## A) Objectives

Objectives of original proposal were met and the results are outlined in this final report.

## B) Status of Effort for Duration of Funding

The focus of this project was to acquire and use computer cluster nodes purchased with AFOSR DURIP funding for the multiscale simulation of ionic liquids and the construction of a novel molecular design methodology. During the funding period atomistic and coarse-grained (CG) molecular dynamics (MD) simulations of ionic liquids were performed to study the structure and dynamical properties of the ionic liquids. An effective force coarse graining (EF-CG) method was developed to build transferable CG models, and a systematic inverse coarse-graining molecular design approach was developed based on the EF-CG model and an empirical CG-structure-property relationship.

## C) Significant Work Accomplished

The behavior of the ionic liquid/vacuum interface was studied by applying the multi-scale coarse-graining (MS-CG)<sup>[2-4]</sup> method,<sup>[5]</sup> and the MSCG models of 1-alkyl-3-methylimidazolium-based ionic liquids (IL) with alkyl substituent of different lengths were developed. When the chain length of cations increases, a monolayer ordering will change to a unique multilayer ordering and the surface tension will decrease and approaches a constant, consistent with Langmuir theory<sup>[6]</sup> and experiments.<sup>[7, 8]</sup> A detailed orientational study demonstrated that the alkyl chain of cations prefers to align parallel to the surface normal while the aromatic ring tends to be perpendicular to the surface normal. The surface electron density profile shows that the surface electron density oscillations are mainly contributed by cations. Figure 1 (a) through (d) (shown at right) illustrates the 1-dodecyl-3-methylimidazolium/NO<sub>3</sub><sup>-</sup> system, revealing a clear multilayer ordering: (a) all CG sites; (b) CG sites of aromatic rings, the CH<sub>3</sub> groups on long side chains and anions; (c) CG sites of the CH<sub>3</sub> groups on long side chains; and (d) CG sites of aromatic rings and anions.

An effective force coarse-graining (EF-CG) method was developed to build transferable coarse-graining force field<sup>[1]</sup>. In the EF-CG method, the effective pairwise forces between coarse-grained sites were calculated by taking the average of the atomistic forces between the corresponding atomic groups in configurations sampled from equilibrated atomistic molecular dynamics simulations. When compared with more general MS-CG method, the EF-CG method keeps the transferable part of the MS-CG at the cost of slightly reduced structural accuracy. Therefore, as an alternative to the




Figure 1<sup>[5]</sup>



MS-CG method, the EF-CG method determines the CG force fields with higher transferability but reduced structural accuracy. The computer hardware purchased with the DURIP funding greatly facilitated the development and testing of the EF-CG method.




Figure 2<sup>[1]</sup>

For ionic liquid systems, the EF-CG method reproduced all liquid state structure with little loss in local structure details. In particular, the 1-ethyl-3-methylimidazolium (EMIM)/NO<sub>3</sub><sup>-</sup> system was studied by using the EF-CG method. Figure 2 (shown above) (a) and (b) are the all-atom and EF-CG models for EMIM/NO<sub>3</sub><sup>-</sup>. Figure 3 below shows the comparison of radial distribution functions (RDFs) for different CG sites between EF-CG and all-atom MD results.




Figure 3<sup>[1]</sup>

The transferability of the EF-CG force field was studied for different ionic liquid structures<sup>[9]</sup> and different temperatures.<sup>[1, 9]</sup> One IL was used to reproduce the structures of IL with different side chain length and the transferability of any two ionic liquids decreased when

increasing the difference between the side chain lengths of that two ionic liquids. Figure 4 (shown below) depicts selected RDFs for the 1-decyl-3-methylimidazolium/ $\text{NO}_3^-$  (C10): In Figure 4 (a) is the RDF for head groups and (b) is the RDF for tail groups. The head group is taken as the aromatic ring of the cations with the  $\text{CH}_3$  on the long alkyl side chain as the tail group. For the black curve, the EF-CG force field was calculated from an all atom simulation of 1-octyl-3-methylimidazolium/ $\text{NO}_3^-$  (C8), and for the red curve, the EF-CG force field was calculated from an all atom simulation of a C10 system. The RDFs show very good agreement between the EF-CG force field coming from C8 and C10. Therefore, the EF-CG force field has good transferability between ILs with different side chain lengths.




Figure 4

The EF-CG simulation, under different temperatures, revealed the EF-CG force fields have very good transferability within the temperature range studied (298K-700K).<sup>[1, 9]</sup> Figure 5 below shows the RDFs of 1-hexyl-3-methylpyridinium/ $\text{PF}_6^-$  (HMPM/ $\text{PF}_6^-$ ) systems at 400K. Figure 5 (a) is for head groups and (b) is for tail groups. For the black curve, the EF-CG force field was calculated from an all-atom simulation of HMPM/ $\text{PF}_6^-$  at 298K and the EF-CG force field for the red curve was calculated from an all-atom MD simulation of HMPM/ $\text{PF}_6^-$  at 400K. The results show the very good temperature transferability of EF-CG force field.




Figure 5

Based on the EF-CG method, an empirical relationship between CG structure and a desired set properties, an inverse coarse-graining (ICG) approach, has been developed with the following procedure:<sup>[9]</sup> 1) several sets of ionic liquids were selected to represent the general properties of a wide range of such liquids; 2) all atom MD simulations were carried out for each selection; 3) EF-CG models were built for each IL and the force fields for each of the CG sites were added to the EF-CG library. At the same time, an empirical CG-structure-property network was built based on the relationship between CG structures and the target properties of density and heterogeneity. Given the desired properties (i.e., density and heterogeneous order parameter<sup>[10]</sup>), the CG-structure-property network provided predicted CG structures. Next, a search of the EF-CG library was conducted to construct a proper force field for the CG structures which came from the CG-structure-property network. Finally, an EF-CG simulation was performed for the constructed CG models to choose the best CG representations followed by an all atom MD simulation with each of the CG representations, giving the best all-atom representations, which provide the final IL result for a given desired density and heterogeneous order parameter.

#### Description of Computational Codes

DL\_POLY: [http://www.csc.scitech.ac.uk/ccg/software/DL\\_POLY/](http://www.csc.scitech.ac.uk/ccg/software/DL_POLY/)

Gromacs: <http://www.gromacs.org/>

Revised DL\_POLY: MD simulation package specially used to do the polarizable model calculation.

MS-CG: computational codes used to build MS-CG model based on all atom simulations

EF-CG: computational codes used to build EF-CG models based on all atom simulations

#### Description of the Computing Cluster Purchased with DURIP Funding

A new 384-core Dell PE 1950 rackmount server cluster, purchased with the awarded DURIP funds (FA9550-01-1-0082), was built and has been used to help fulfill the computational requirements of the research projects outlined above. The DURIP funds used to purchase this cluster equipment which included 48 compute nodes, 3 service/management nodes, a fully non-blocking Qlogic SDR InfiniBand (IB) switch, a Gigabit Ethernet (GigE) switch, and an 8-TB global scratch space. Each compute node has two 2.33/2.66 GHz quad-core Intel Xeon processors (for a total of 8 cores per node), 16 GB RAM (2 GB per core), 500 GB local disk, and an x4 SDR IB Qlogic/Pathscale "InfiniPath" single-port HCA (Host Channel Adapter) card over x8 PCI-E for very low-latency and high-bandwidth communication.

#### References:

1. Wang, Y., et al., "Effective Force Coarse Graining " In Preparation 2008.
2. Izvekov, S. and G.A. Voth, "A multiscale coarse-graining method for biomolecular systems." *Journal of Physical Chemistry B*, 2005. **109**(7): p. 2469-2473.
3. Izvekov, S. and G.A. Voth, "Multiscale coarse graining of liquid-state systems." *Journal of Chemical Physics*, 2005. **123**(13): p. 134105.
4. Wang, Y.T., et al., "Multiscale coarse-graining of ionic liquids." *Journal of Physical Chemistry B*, 2006. **110**(8): p. 3564-3575.
5. Jiang, W., et al., "A multiscale coarse-graining study of the liquid/vacuum interface of room-temperature ionic liquids with alkyl substituents of different lengths." *Journal of Physical Chemistry C*, 2008. **112**(4): p. 1132-1139.

Comprehensive Final Report: Gregory A. Voth, PI  
Contract/Grant title: Multiscale Simulation of High Energy Density Ionic Liquids  
Contract/Grant#: FA9550-01-1-0082  
Reporting Period: Total Award Period: 11/18/2003 – 12/31/2006

6. Langmuir, I., Colloid Symp. Monogr. , 1925: p. 48.
7. Law, G. and P.R. Watson, "*Surface tension measurements of N-alkylimidazolium ionic liquids.*" Langmuir, 2001. **17**(20): p. 6138-6141.
8. Mach, P., et al., "*Surface tension obtained from various smectic free-standing films: The molecular origin of surface tension.*" Langmuir, 1998. **14**(15): p. 4330-4341.
9. Feng, S., Y. Wang, and G.A. Voth, "*Inverse Coarse-graining : A New Tool for Molecular Design.*" In Preparation, 2008.
10. Wang, Y.T. and G.A. Voth, "*Tail aggregation and domain diffusion in ionic liquids.*" Journal of Physical Chemistry B, 2006. **110**(37): p. 18601-18608.